

Central limit theorems for nonlinear hierarchical sequences of random variables

Jung M. Woo and Jan Wehr
University of Arizona

July 10, 1999

Abstract

We study central limit theorems for certain nonlinear sequences of random variables. In particular, we prove the central limit theorems for the bounded conductivity of the random resistor networks on hierarchical lattices

Key words and phrases:

AMS 1991 subject classification number

Running head:

1 Introduction

Let D be a closed interval of $\mathbf{R}$ (bounded or unbounded) and $f : D^k \rightarrow D$ ($k \geq 2$) be a continuous function. Let x_0 be a D -valued random variable with a distribution μ_0 . We define a hierarchical sequence of random variables $\{x_n : n = 0, 1, 2, \dots\}$ inductively by

$$x_n = f(x_{n-1,1}, x_{n-1,2}, \dots, x_{n-1,k}) \quad (1)$$

where for each $n \geq 1$ $x_{n-1,1} x_{n-1,2} \dots, x_{n-1,k}$ are independent identically distributed (shortly IID) random variables having the same distribution as x_{n-1} . For each $n \geq 0$ let us denote a distribution of x_n by μ_n . We thus write

$$\begin{aligned} x_n &= \underbrace{f \circ f \circ \dots \circ f}_{n \text{ times}}(x_0) \\ \mu_n &= \underbrace{f \circ f \circ \dots \circ f}_{n \text{ times}}(\mu_0) \end{aligned} \quad (2)$$

We are interested in central limit theorem for the hierarchical sequence x_n under suitable assumptions of f . Assume that f is twice continuously differentiable at $(c_n, c_n, \dots, c_n)$ where $c_n = E[x_n]$. Let

$$\alpha_{n,i} = \frac{\partial f}{\partial u_i}(c_n, c_n, \dots, c_n)$$

By expanding x_{n+1} at $(c_n, c_n, \dots, c_n)$, we get

$$x_{n+1} = f(c_n, c_n, \dots, c_n) + \sum_{i=1}^k \alpha_{n,i}(x_{n,i} - c_n) + \text{second order} + \dots \quad (3)$$

For each n let

$$d_n = f(c_n, c_n, \dots, c_n) - c_n \sum_{i=1}^k \alpha_{n,i}$$

and

$$z_n = x_{n+1} - \sum_{i=1}^k \alpha_{n,i} x_{n,i} \quad (4)$$

Then

$$z_n = d_n + \text{second order} + \dots$$

Hence x_{n+1} can be viewed as a linear function of $x_{n,i}$ plus some random variable z_n of “small” variance. The following central limit theorem applies to sequences of this kind (not necessarily defined by iterations).

Theorem 1 For all $n \geq 0$, let k_n be a positive integer greater than or equal to 2. Assume that a sequence of real-valued random variables $\{x_n, n = 0, 1, 2, \dots\}$ satisfies the following recursive relation.

$$x_{n+1} = \sum_{i=1}^{k_n} \alpha_{n,i} x_{n,i} + z_n \quad (5)$$

where for each $n \geq 0$ $\alpha_{n,i} (i \leq k_n)$ are real numbers, z_n is a real-valued random variable with $E[z_n^2] < \infty$, and $\{x_{n,i} : i = 1, \dots, k_n\}$ are IID random variables with same distribution as x_n . We also assume that $E[x_0^2] < \infty$ and that there exists a $\delta > 0$ such that for all n $|\alpha_{n,i}| \geq \delta$ and $|\alpha_{n,j}| \geq \delta$ for at least two distinct indices i and j (“mixing” property). Let $\lambda_n = \sqrt{\sum_{i=1}^{k_n} \alpha_{n,i}^2}$ and $\sup_n \lambda_n < \infty$. Furthermore, assume that there exist $\delta_1 > 0$, $\delta_2 > 0$, $C_1 > 0$ and $C_2 > 0$ with $\delta_1 < \delta_2$ such that for any n

$$\begin{aligned} \text{Var}[x_n] &\geq C_1^2 \lambda_0^2 \lambda_1^2 \dots \lambda_{n-1}^2 (1 - \delta_1)^{2n} \\ \text{Var}[z_n] &\leq C_2^2 \lambda_0^2 \lambda_1^2 \lambda_2^2 \dots \lambda_{n-1}^2 (1 - \delta_2)^{2n} \end{aligned} \quad (6)$$

Then $\frac{x_n - E x_n}{\sqrt{\text{Var}[x_n]}}$ converges in distribution to a unit normal variable.

We will prove Theorem 1 in section 2. To apply Theorem 1 to some hierarchical sequence of real-valued random variables x_n , we have to prove the variance bounds (6). In order to prove the variance bounds in many cases, we need weak laws of large numbers and large deviation estimates. In section 3, we will prove weak laws of large numbers and large deviation estimates for some class of functions f .

We will say that a continuous function $f : D^k \rightarrow D$ (where D is a closed interval) is averaging if the following three conditions hold:

1. For all $u_i \in D$ ($i = 1, 2, \dots, k$)

$$\min_i u_i \leq f(u_1, u_2, \dots, u_k) \leq \max_i u_i \quad (7)$$

2. f is monotone, that is, for all u_i and u'_i (with $u_i \leq u'_i$)

$$f(u_1, u_2, \dots, u_k) \leq f(u'_1, u'_2, \dots, u'_k) \quad (8)$$

3. if equality holds in 2 then

$$u_i = u'_i \text{ and } u_j = u'_j \text{ for at least two distinct indices } i \text{ and } j \quad (9)$$

The following Theorem 2, proved in section 4, is one of the applications of Theorem 1.

Theorem 2 *Let $[a, b]$ be a closed and bounded interval with $a < b$. Let $f : [a, b]^k \rightarrow [a, b]$ be averaging. Assume that the essential range of $[a, b]$ -valued random variable x_0*

$$\mathcal{R}(x_0) = \{u \in [a, b] : \text{for any } \epsilon > 0 \mathbf{P}[|x_0 - u| > \epsilon] > 0\}$$

is connected and consists of more than one point. Suppose there exists $c \in (a, b)$ such that x_n converges to c in probability (a weak law of large numbers). Also assume that there exists $\epsilon_1 > 0$ such that $(c - \epsilon_1, c + \epsilon_1) \subset (a, b)$ and f is twice continuously differentiable on $(c - \epsilon_1, c + \epsilon_1)^k$ and that $\frac{\partial f}{\partial u_i}(c, c, \dots, c) > 0$ for two distinct indices i . Then

$$\frac{x_n - E[x_n]}{\sqrt{\text{Var}[x_n]}}$$

converges to a unit normal variable in distribution.

Another application of Theorem 1 is to prove a central limit theorem for the conductivity of random resistor networks on hierarchical lattices. Let us define hierarchical lattices first. A detail description of various hierarchical lattices can be found in [3] and [4]. Let $\mathbf{G} = (\mathcal{S}, \mathcal{B}, (s_t, s_b))$ be a graph with site set $\mathcal{S}$, bond set $\mathcal{B}$, top site s_t and bottom site s_b . The top site and bottom site are called surface sites since they are the sites where we apply the potential difference. All other sites are called internal sites. Let k be the total number of bonds of $\mathbf{G}$. We define a hierarchical lattice $\mathbf{H} = \{\mathbf{H}_n : n = 0, 1, 2\}$ (constructed from the fixed graph $\mathbf{G}$) inductively. At level 0, H_0 is a single bond. At level n , a graph $\mathbf{H}_n$ is constructed from $\mathbf{H}_{n-1}$ by replacing each bond of $\mathbf{H}_{n-1}$ by the fixed graph $\mathbf{G}$. Since at each level each bond is replaced by the graph $\mathbf{G}$ having k bonds, $\mathbf{H}_n$ consists of total k^n bonds, many internal sites and two surface sites where we apply the potential difference. We will assume that the graph $\mathbf{G}$ used in constructing the hierarchical lattice $\mathbf{H}_n$, is connected and there exist at least two bond-disjoint self-avoiding paths connecting two surface sites of $\mathbf{G}$. we will also assume that for any bond b of $\mathbf{G}$ there exists a self-avoiding path passing through the bond b and connecting two surface sites. Furthermore we will assume any self-avoiding path connecting two surface sites of $\mathbf{G}$ has at least length 2. Let us order all bonds of $\mathbf{G}$ in a specific way. For each $i = 1, \dots, k$ let $u_i \geq 0$ be the conductance of the i -th bond of $\mathbf{G}$. Let $h(u_1, u_2, \dots, u_k)$ be the conductance

of the graph $\mathbf{G}$. Define $f : [0, \infty)^k \rightarrow [0, \infty)$ by

$$f(u_1, u_2, \dots, u_k) = \frac{h(u_1, u_2, \dots, u_k)}{h(1, 1, \dots, 1)} \quad (10)$$

Then this function f is nonnegative continuous, monotone increasing, homogeneous of degree one, concave, and $f(1, 1, \dots, 1) = 1$ (See [5]). Let $x_{0,i}$ be IID random variables with a common distribution μ_0 where $x_{0,i}$ represents the random conductance of i -th bond of $\mathbf{G}$. Then the random conductance $\hat{x}_n$ of the hierarchical lattice $\mathbf{H}_n$ is given by

$$\hat{x}_n = \underbrace{h \circ h \circ \dots \circ h}_{n \text{ times}}(x_0) \quad (11)$$

Clearly a normalized random conductance $x_n = h(1, 1, \dots, 1)^{-n} \hat{x}_n$ is

$$x_n = \underbrace{f \circ f \circ \dots \circ f}_{n \text{ times}}(x_0) \quad (12)$$

Let x_0 be a bounded nonnegative random variable. Let

$$p = \mathbf{P}[x_0 > 0]$$

and

$$g(p) = \mathbf{P}[x_1 > 0]$$

Note that the function $g(p)$ does not depend on the distribution of x_0 but depends only on the parameter p . In [5], Shneiberg proved that there exists $c(\mu_0)$ such that the normalized conductivity x_n converges to $c(\mu_0)$ in probability and that there exists a unique fixed point $p_c(\mathbf{G}) \in (0, 1)$ of g (that is $g(p) = p$) such that

$$c(\mu_0) = \begin{cases} 0 & \text{if } p \leq p_c \\ \text{positive} & \text{if } p > p_c \end{cases} \quad (13)$$

The following Theorem 3 is a central limit theorem for the conductivity of random resistor networks on hierarchical lattices. The proof will be given in section 4.

Theorem 3 *Let $\hat{x}_n$ be the conductivity of a random resistor network on a hierarchical lattice $\mathbf{H}$ constructed using a fixed graph $\mathbf{G}$ described above. Assume that a random variable x_0 is nonnegative, bounded and not almost surely*

a constant. Also let $\mathbf{P}[x_0 > 0] > p_c(\mathbf{G})$. where $p_c(\mathbf{G})$ is a unique fixed point of g mentioned above. Then the random variables

$$\frac{\hat{x}_n - \mathbf{E}[\hat{x}_n]}{\sqrt{\mathbf{Var}[\hat{x}_n]}}$$

converge in distribution to a unit normal variable.

Remark: For $\mathbf{P}[x_0 > 0] \leq p_c(\mathbf{G})$ we expect a non-Gaussian behavior for $\hat{x}_n$.

2 Proof of Theorem 1

In this section we use the notations introduced in Theorem 1 as well as the following. For all $n \geq 0$ let

$$y_n = \sum_{i=1}^{k_n} \alpha_{n,i} x_{n,i} \quad (14)$$

and for all $n \geq 1$ let

$$\begin{aligned} \widetilde{x}_n &= \frac{x_n - \mathbf{E}[x_n]}{\lambda_0 \lambda_1 \cdots \lambda_{n-1}} \\ \widetilde{y}_n &= \frac{y_n - \mathbf{E}[y_n]}{\lambda_0 \lambda_1 \cdots \lambda_{n-1}} \\ \widetilde{z}_n &= \frac{z_n - \mathbf{E}[z_n]}{\lambda_0 \lambda_1 \cdots \lambda_{n-1}} \end{aligned} \quad (15)$$

Then the assumptions in Theorem 1 say that for all n

$$\begin{aligned} \widetilde{x}_{n+1} &= \frac{1}{\lambda_n} (\widetilde{y}_n + \widetilde{z}_n) \\ \sqrt{\mathbf{Var}[\widetilde{y}_n]} &= \lambda_n \sqrt{\mathbf{Var}[\widetilde{x}_n]} \\ \sqrt{\mathbf{Var}[\widetilde{x}_n]} &\geq C_1 (1 - \delta_1)^n \\ \sqrt{\mathbf{Var}[\widetilde{z}_n]} &\leq C_2 (1 - \delta_2)^n \end{aligned} \quad (16)$$

Lemma 1 *Let*

$$\sigma_\infty = \lim_{n \rightarrow \infty} \sqrt{\mathbf{Var}[\widetilde{x}_n]}$$

Then σ_∞ exists and is a positive real number. Moreover there exist positive real constants C_3 , C_4 , and C_5 such that for all n

$$\begin{aligned} \inf_n \sqrt{\mathbf{Var}[\widetilde{x}_n]} &\geq C_3 \\ \sup_n \sqrt{\mathbf{Var}[\widetilde{x}_n]} &\leq C_4 \\ \sup_n \sqrt{\mathbf{Var}[\widetilde{y}_n]} &\leq C_5 \end{aligned} \tag{17}$$

Proof: From (16) and by the triangle inequality,

$$\begin{aligned} &\left| \sqrt{\mathbf{Var}[\widetilde{x}_{n+1}]} - \sqrt{\mathbf{Var}[\widetilde{x}_n]} \right| \\ &= \left| \sqrt{\mathbf{Var}[\widetilde{x}_{n+1}]} - \frac{1}{\lambda_n} \sqrt{\mathbf{Var}[\widetilde{y}_n]} \right| \\ &\leq \frac{1}{\lambda_n} \sqrt{\mathbf{Var}[\widetilde{z}_n]} \leq \frac{1}{\underline{\lambda}} C_2 (1 - \delta_2)^n \end{aligned}$$

where $\underline{\lambda} = \inf_n \lambda_n > \delta$. By summing up the above inequality for all $n \geq m$, we can find a positive constant C_6 such that for all m

$$\sum_{n=m}^{\infty} \left| \sqrt{\mathbf{Var}[\widetilde{x}_{n+1}]} - \sqrt{\mathbf{Var}[\widetilde{x}_n]} \right| \leq C_6 (1 - \delta_2)^m$$

Since the above series converges absolutely, in particular σ_∞ exists (and is finite) and for all m

$$\left| \sigma_\infty - \sqrt{\mathbf{Var}[\widetilde{x}_m]} \right| \leq C_6 (1 - \delta_2)^m$$

Therefore from (16) and from the above inequality, for all m we have

$$C_1 (1 - \delta_1)^m \leq \sqrt{\mathbf{Var}[\widetilde{x}_m]} \leq \sigma_\infty + C_6 (1 - \delta_2)^m \tag{18}$$

Since the above inequality (18) is true for all m , from $\delta_2 > \delta_1$ σ_∞ cannot be zero. Three inequalities of Lemma 1 follow immediately from the positivity of σ_∞ , and (16).

Lemma 2 *Let C_7 be any positive constant, and let X and Y be random variables with zero means and variances less than C_7^2 . Then for all t ($|t| < \frac{1}{C_7}$)*

$$|\ln \mathbf{E}[\exp(itX)] - \ln \mathbf{E}[\exp(itY)]| \leq 4C_7 t^2 \sqrt{\mathbf{E}[(X - Y)^2]} \quad (19)$$

Also we have

$$\lim_{t \rightarrow 0} \frac{\ln \mathbf{E}[\exp(itX)]}{t^2} = -\frac{1}{2} \mathbf{Var}[X] \quad (20)$$

Proof: Since $|e^{iu} - 1 - iu| \leq \frac{u^2}{2}$ for any real number u ,

$$|\mathbf{E}[\exp(itX)] - 1| \leq t^2 \frac{\mathbf{E}[X^2]}{2}$$

Therefore for $|t| < \frac{1}{C_7}$, the value $\mathbf{E}[\exp(itX)]$ lies inside of a circle of radius $1/2$ centered at $(1, 0)$ in the complex plane (similarly to Y). Hence for $|t| < \frac{1}{C_7}$ the logarithmic function $\ln \mathbf{E}[\exp(itX)]$ is single-valued. Since $|\ln u - \ln v| \leq 2|u - v|$ for any complex number u and v with $|u - 1| \leq \frac{1}{2}$ and $|v - 1| \leq \frac{1}{2}$, for $|t| < \frac{1}{C_7}$ we have

$$\begin{aligned} & |\ln \mathbf{E}[\exp(itX)] - \ln \mathbf{E}[\exp(itY)]| \\ & \leq 2 |\mathbf{E}[\exp(itX)] - \mathbf{E}[\exp(itY)]| = 2 |\mathbf{E}[\exp(itX)(1 - \exp(it(Y - X)))]| \\ & \leq 2 \mathbf{E}[|\exp(itX) - 1| |1 - \exp(it(Y - X))|] + 2 \mathbf{E}[|1 - \exp(it(Y - X))|] \\ & \leq 2t^2 \mathbf{E}[|X||Y - X|] + t^2 \mathbf{E}[(Y - X)^2] \\ & \leq 4t^2 C_7 \sqrt{\mathbf{E}[(X - Y)^2]} \end{aligned}$$

where in the last part we used Cauchy-Schwarz inequality and the triangle inequality. The second part of Lemma 2 is elementary and the proof can be found in many basic probability textbooks (for example see [2]).

For all n let us define a characteristic function ϕ_n of the normalized random variable $\widetilde{x}_n$ by

$$\phi_n(t) = \mathbf{E}[\exp(it\widetilde{x}_n)]$$

Lemma 3 *There exists a positive constant C_8 such that for any $|t| < \frac{1}{C_8}$*

$$\lim_{n \rightarrow \infty} \ln \phi_n(t) = -\frac{1}{2} \sigma_\infty^2 t^2$$

(the convergence is uniform on $|t| < \frac{1}{C_8}$). Therefore $\widetilde{x}_n$ converges in distribution to a normal variable with mean zero and variance σ_∞^2 .

Proof: From Lemma 1 and Lemma 2, it follows that there exists a positive constant C_8 such that for any $|t| < \frac{1}{C_8}$ and any n

$$\left| \ln \mathbf{E}[\exp(it\widetilde{x_{n+1}})] - \ln \mathbf{E}[\exp(it\frac{1}{\lambda_n}\widetilde{y_n})] \right| \leq 4C_8t^2 \sqrt{\mathbf{Var} \left[\frac{\widetilde{z_n}}{\lambda_n} \right]}$$

For all n and $i \leq k_n$, let $\beta_{n,i} = \frac{\alpha_{n,i}}{\lambda_n}$. Then from the above inequality and from (16), there is a positive constant C_9 such that for all $|t| \leq \frac{1}{C_8}$

$$\left| \ln \phi_{n+1}(t) - \sum_{i=1}^{k_n} \ln \phi_n(\beta_{n,i}t) \right| \leq C_9t^2(1 - \delta_2)^n \quad (21)$$

Note that for any real $\gamma_{n,j}$ satisfying $\sum_j \gamma_{n,j}^2 = 1$, we have

$$\sum_j \left| \ln \phi_{n+1}(\gamma_{n,j}t) - \sum_{i=1}^{k_n} \ln \phi_n(\gamma_{n,j}\beta_{n,i}t) \right| \leq C_9t^2(1 - \delta_2)^n \quad (22)$$

From the above inequality (22), it follows that for all $|t| \leq \frac{1}{C_8}$

$$\begin{aligned} & \left| \ln \phi_{n+m}(t) - \sum_{i_1, i_2, \dots, i_m} \ln \phi_n(t\beta_{n,i_1}\beta_{n+1,i_2} \cdots \beta_{n+m-1,i_m}) \right| \\ & \leq C_9t^2 \sum_{p=0}^{m-1} (1 - \delta_2)^{n+p} \leq \frac{C_9}{\delta_2} t^2 (1 - \delta_2)^n \end{aligned} \quad (23)$$

Note that by one of the assumptions in Theorem 1 (“mixing” property)

$$\max_{i_1, i_2, \dots, i_m} \beta_{n,i_1}\beta_{n+1,i_2} \cdots \beta_{n+m-1,i_m}$$

converges to zero when $m \rightarrow \infty$. From the second part of Lemma 2, it follows that for any n and any $\epsilon_1 > 0$ there exists an $M_1(n, \epsilon_1)$ such that for all $m \geq M_1$

$$\begin{aligned} & \left| \ln \phi_n(t\beta_{n,i_1}\beta_{n+1,i_2} \cdots \beta_{n+m-1,i_m}) + \frac{1}{2}t^2(\beta_{n,i_1}\beta_{n+1,i_2} \cdots \beta_{n+m-1,i_m})^2 \mathbf{Var}[\widetilde{x_n}] \right| \\ & \leq \epsilon_1 t^2 (\beta_{n,i_1}\beta_{n+1,i_2} \cdots \beta_{n+m-1,i_m})^2 \end{aligned}$$

Hence for any $m \geq M_1$, by summing the above inequality over all $i_1, i_2, \dots, i_m$, we obtain

$$\left| \sum_{i_1, i_2, \dots, i_m} \ln \phi_n(t\beta_{n, i_1} \beta_{n+1, i_2} \dots \beta_{n+m-1, i_m}) + \frac{1}{2} t^2 \mathbf{Var}[\widetilde{x}_n] \right| \leq \epsilon_1 t^2 \quad (24)$$

From (23) and (24), using the triangle inequality, we have for all n , $\epsilon_1 > 0$, and $|t| < \frac{1}{C_8}$

$$\limsup_{m \rightarrow \infty} \left| \ln \phi_{n+m}(t) + \frac{1}{2} t^2 \mathbf{Var}[\widetilde{x}_n] \right| \leq \epsilon_1 t^2 + \frac{C_9}{\delta_2} t^2 (1 - \delta_2)^n$$

Since $\epsilon_1 > 0$ is arbitrary, the proof is finished by taking the limit $n \rightarrow \infty$.

Proof of Theorem 1. From Lemma 3, $\frac{\widetilde{x}_n}{\sqrt{\mathbf{Var}[\widetilde{x}_n]}}$ converges to a unit normal variable in distribution. Since

$$\frac{x_n - \mathbf{E}[x_n]}{\sqrt{\mathbf{Var}[x_n]}} = \frac{\widetilde{x}_n}{\sqrt{\mathbf{Var}[\widetilde{x}_n]}}$$

we finish the proof of Theorem 1.

3 Law of large numbers

In this section we state and prove weak law of large numbers and large deviation estimate used later to prove central limit theorems for hierarchical sequences of real-valued random variables. To be more precise, we define some terminology below. A sequence of real-valued random variables x_n satisfies a weak law of large numbers if there exists a real number c such that for any $\epsilon > 0$

$$\lim_{n \rightarrow \infty} \mathbf{P}[|x_n - c| > \epsilon] = 0 \quad (25)$$

We say that x_n satisfies a large deviation estimate if there exists a real number c such that for any $\epsilon_1 > 0$ and $\epsilon_2 > 0$, there exists $M > 0$ such that for all n

$$\mathbf{P}[|x_n - c| > \epsilon_1] \leq M \epsilon_2^n \quad (26)$$

The above is not the strongest large deviation estimate satisfied by hierarchical sequences studied in this paper, but it is a sufficient condition for central limit theorems.

Proposition 4 *Let D be a closed interval (bounded or unbounded) of $\mathbf{R}$. Let $f : D^k \rightarrow D$ be differentiable and x_0 be a D -valued random variable with $\mathbf{E}[\exp(\delta|x_0|)] < \infty$ for some $\delta > 0$. Also assume that there exists $\epsilon > 0$ such that*

$$\sup_{(u_1, \dots, u_k)} \sum_{i=1}^k \left| \frac{\partial f}{\partial u_i}(u_1, u_2, \dots, u_k) \right| < 1 - \epsilon$$

Then a weak law of large numbers and a large deviation estimate hold.

Proof: Let us define $g(u) = f(u, u, \dots, u)$. Then by the mean value theorem $|g(u) - g(v)| \leq (1 - \epsilon)|u - v|$. Thus g is a contraction mapping on D and, consequently has a unique fixed point $c \in D$, that is $f(c, c, \dots, c) = c$. Then by the mean value theorem and our assumptions on f ,

$$|x_{n+1} - c| = |f(x_{n,1}, x_{n,2}, \dots, x_{n,k}) - f(c, c, \dots, c)| \leq (1 - \epsilon) \max_{i \in \{1, \dots, k\}} |x_{n,i} - c|$$

Therefore for any positive real number v

$$\mathbf{P}[|x_{n+1} - c| > v] \leq k \mathbf{P}[|x_n - c| > \frac{v}{1 - \epsilon}]$$

From the above inequality, for any $v > 0$ and for all n we have

$$\mathbf{P}[|x_n - c| > v] \leq k^n \mathbf{P}[|x_0 - c| > \frac{v}{(1 - \epsilon)^n}] \leq k^n \mathbf{E}[\exp(\delta(|x_0 - c| - \frac{v}{(1 - \epsilon)^n}))]$$

where in the last equation we used Chebyshev's inequality. Weak law of large numbers and large deviation estimate follows immediately from the above inequality.

Proposition 5 *Let $[a, b]$ be a closed and bounded interval of $\mathbf{R}$ with $a < b$, and $f : [a, b]^k \rightarrow [a, b]$ be averaging. Also let x_0 be a $[a, b]$ -valued random variable. Furthermore assume that there exists $c \in [a, b]$ such that x_n converges to c in probability (weak law of large numbers). Then a large deviation estimate holds for the same constant c .*

Proof: By the symmetry of definitions, it is enough to show that for any $\epsilon > 0$ and for any $u < c$ there exists $M > 0$ such that for all n

$$\mathbf{P}[x_n < u] < M\epsilon^n \tag{27}$$

In case $c = a$, the above inequality is obvious, hence we will assume that $c \in (a, b]$. Let $F_n(u) = \mathbf{P}[x_n \leq u]$ be the distribution function of x_n and let

$$Q = \{u \in [a, b] : \text{for any } \epsilon > 0 \text{ there exists an } M \text{ such that } F_n(u) \leq M\epsilon^n\} \quad (28)$$

We only need to prove that $\sup Q = c$. Without loss of generality we will assume that f is symmetric under the exchange of k variables. First we will show that for some positive ϵ_1 , $a + \epsilon_1 \in Q$. Since $f(a, c, c, \dots, c) > a$ (by monotonicity) there exists $\epsilon_1 \in (0, c - a)$ such that

$$f(a, c - \epsilon_1, c - \epsilon_1, \dots, c - \epsilon_1) > a + \epsilon_1$$

For such ϵ_1

$$\begin{aligned} & \mathbf{P}[x_{n+1} \leq a + \epsilon_1] \\ & \leq k! \mathbf{P}[f(x_{n,1}, x_{n,2}, \dots, x_{n,k}) \leq a + \epsilon_1; x_{n,1} \leq x_{n,2} \leq \dots \leq x_{n,k}; x_{n,1} \leq a + \epsilon_1] \\ & \leq k! \mathbf{P}[x_{n,1} \leq a + \epsilon_1; f(a, x_{n,2}, x_{n,2}, \dots, x_{n,2}) \leq a + \epsilon_1] \\ & \leq k! \mathbf{P}[x_{n,1} \leq a + \epsilon_1; x_{n,2} \leq c - \epsilon_1] \\ & \leq k! \mathbf{P}[x_n \leq c - \epsilon_1] \mathbf{P}[x_n \leq a + \epsilon_1] \end{aligned}$$

From the above inequality and by the weak law of large numbers, it follows that $a + \epsilon_1 \in Q$. Now we will prove that $\sup Q = c$. We will prove this by contradiction. Let $u = \sup Q < c$. First note that $u \geq a + \epsilon_1$. Since $f(u, c, c, \dots, c) > u$ (by monotonicity) there exists $\epsilon_2 \in (0, \epsilon_1)$ such that

$$f(u - \epsilon_2, c - \epsilon_2, c - \epsilon_2, \dots, c - \epsilon_2) > u + \epsilon_2$$

For such ϵ_2

$$\begin{aligned} & \mathbf{P}[x_{n+1} \leq u + \epsilon_2] \\ & \leq k! \mathbf{P}[f(x_{n,1}, x_{n,2}, \dots, x_{n,k}) \leq u + \epsilon_2; x_{n,1} \leq x_{n,2} \leq \dots \leq x_{n,k}; x_{n,1} \leq u + \epsilon_2] \\ & \leq k! \mathbf{P}[x_{n,1} \leq u - \epsilon_2] + \\ & \quad k! \mathbf{P}[u - \epsilon_2 \leq x_{n,1} \leq u + \epsilon_2; f(u - \epsilon_2, x_{n,2}, x_{n,2}, \dots, x_{n,2}) \leq u + \epsilon_2] \\ & \leq k! \mathbf{P}[x_{n,1} \leq u - \epsilon_2] + k! \mathbf{P}[x_{n,1} \leq u + \epsilon_2; x_{n,2} \leq c - \epsilon_2] \\ & \leq k! \mathbf{P}[x_n \leq u - \epsilon_2] + k! \mathbf{P}[x_n < c - \epsilon_2] \mathbf{P}[x_n \leq u + \epsilon_2] \end{aligned}$$

Take any $\epsilon > 0$. By weak law of large numbers and from $u - \epsilon_2 \in Q$ (by the assumption of u), there exist N and $M > 0$ such that for all $n \geq N$

$$\mathbf{P}[x_{n+1} \leq u + \epsilon_2] \leq M\left(\frac{\epsilon}{2}\right)^n + \frac{\epsilon}{2} \mathbf{P}[x_n \leq u + \epsilon_2]$$

Obviously by induction that there exists $K > 0$ such that for all n $F_n(u + \epsilon_2) \leq K\epsilon^n$, which is a contradiction by definition of Q . Hence we finished the proof of Proposition 5.

4 Variance bounds

In this section we will prove Theorem 2 and Theorem 3. Let $[a, b]$ be a closed and bounded interval of $\mathbf{R}$. Throughout this section, we assume that $f : [a, b]^k \rightarrow [a, b]$ is averaging and x_0 is $[a, b]$ -valued random variable. We also assume that there exists a real number $c \in (a, b)$ such that x_n , defined as in (1), converges to c in probability. In section 3, we showed for such a function f , our sequence x_n satisfies a large deviation estimate. Furthermore We assume that there exists $\epsilon_1 > 0$ such that f is twice continuously differentiable on $(c - \epsilon_1, c + \epsilon_1)^k$. We use the same notations introduced in section 1, section 2 and the following as well. Let

$$\alpha_i = \frac{\partial f}{\partial u_i}(c, c, \dots, c)$$

$$\lambda = \sqrt{\sum_{i=1}^k \alpha_i^2}$$

$$\underline{\lambda} = \inf_n \lambda_n$$

Let a random variable x'_0 be an independent copy of x_0 , and for all n define x'_n , y'_n and z'_n similarly as we defined x_n , y_n , and z_n in (2),(4) and (14). We have the following variance bounds.

Proposition 6 *For any $\epsilon > 0$ there exists $M > 0$ such that for all n*

$$\mathbf{Var}[z_n] \leq M(\lambda + \epsilon)^{4n}$$

Proof: Since x_n , taking values in a bounded interval $[a, b]$, converges to c in probability, c_n converges to c . Note that there exists $\epsilon_1 > 0$ such that f is twice continuously differentiable on $(c - \epsilon_1, c + \epsilon_1)^k$. Hence for such $\epsilon_1 > 0$ there exists N_1 such that for all $n \geq N_1$ $c - \epsilon_1 < c_n < c + \epsilon_1$. Therefore there exists $M_1 > 0$ such that for all $n \geq N_1$

$$\mathbf{Var}[z_n] \leq M_1 \mathbf{E}[(x_n - c_n)^4] \tag{29}$$

Since for any $\epsilon_2 > 0$ and for all $n \geq N_1$

$$\mathbf{Var}[z_n] \leq M_1 \epsilon_2^2 \mathbf{E}[(x_n - c_n)^2; |x_n - c_n| \leq \epsilon_2] + M_1 \mathbf{E}[(x_n - c_n)^4; |x_n - c_n| \geq \epsilon_2],$$

applying a large deviation estimate to the above inequality, for any ϵ_3 there exists $N_2 > N_1$ such that for all $n \geq N_2$

$$\mathbf{Var}[z_n] \leq \epsilon_3^2 \mathbf{Var}[x_n] + \epsilon_3^{2n}$$

From the triangle inequality and the above inequality, for all $n \geq N_2$ we have

$$\sqrt{\mathbf{Var}[x_{n+1}]} \leq \lambda_n \sqrt{\mathbf{Var}[x_n]} + \sqrt{\mathbf{Var}[z_n]} \leq (\lambda_n + \epsilon_3) \sqrt{\mathbf{Var}[x_n]} + \epsilon_3^n$$

Hence, by the induction on n , for any $\epsilon_3 > 0$ there exists $M_2 > 0$ such that for all n

$$\mathbf{Var}[x_n] \leq M_2 (\lambda + 2\epsilon_3)^{2n} \quad (30)$$

Also note that

$$\mathbf{E}[(x_n - x'_n)^4] = 2\mathbf{E}[(x_n - c_n)^4] + 6(\mathbf{E}[(x_n - c_n)^2])^2$$

Therefore from (29) there exists $M_3 > 0$ and $N_1 > 0$ such that for all $n \geq N_1$

$$\mathbf{Var}[z_n] \leq M_3 \mathbf{E}[(x_n - x'_n)^4] \quad (31)$$

From (14)

$$\begin{aligned} \mathbf{E}[(y_n - y'_n)^4] &\leq \left(\sum_{i=1}^k \alpha_{n,i}^4 \right) \mathbf{E}[(x_n - x'_n)^4] + \lambda_n^4 (\mathbf{E}[(x_n - x'_n)^2])^2 \\ &\leq \lambda_n^4 \left(\mathbf{E}[(x_n - x'_n)^4] + (\mathbf{E}[(x_n - x'_n)^2])^2 \right) \end{aligned}$$

Also since there exists $M_4 > 0$ such that $(a + b)^4 \leq M_4(a^4 + b^4)$ for all real values a and b , and by (4) there exists $M_5 > 0$ such that

$$\mathbf{E}[(z_n - z'_n)^4] \leq M_4 (\mathbf{E}[(z_n - d_n)^4] + \mathbf{E}[(z'_n - d_n)^4]) \leq M_5 \mathbf{E}[(x_n - c_n)^8]$$

Since for any $\epsilon_4 > 0$ there exists M_6 such that for all a and b $(a + b)^4 \leq (1 + \epsilon_4)a^4 + M_6 b^4$, for such ϵ_4 there exists M_7 such that

$$\begin{aligned} \mathbf{E}[(x_{n+1} - x'_{n+1})^4] &= \mathbf{E}[(y_n - y'_n + z_n - z'_n)^4] \\ &\leq (1 + \epsilon_4) \mathbf{E}[(y_n - y'_n)^4] + M_6 \mathbf{E}[(z_n - z'_n)^4] \\ &\leq (1 + \epsilon_4) \lambda_n^4 \left(\mathbf{E}[(x_n - x'_n)^4] + (\mathbf{E}[(x_n - x'_n)^2])^2 \right) + M_7 \mathbf{E}[(x_n - c_n)^8] \end{aligned} \quad (32)$$

Since, by a large deviation estimate, we can show that for any $\epsilon_5 > 0$ there exists $N_3 > N_2$ such that for all $n \geq N_3$

$$M_7 E[(x_n - c_n)^8] \leq \epsilon_5 E[(x_n - c_n)^4] + \epsilon_5^n$$

From the above two inequalities, for any $\epsilon > 0$ there exists an $N_3 > 0$ such that for all $n \geq N_3$

$$\mathbf{E}[(x_{n+1} - x'_{n+1})^4] \leq (\lambda + \epsilon)^4 (\mathbf{E}[(x_n - x'_n)^4] + \mathbf{E}[(x_n - x'_n)^2])^2 + \epsilon^n$$

Hence from (29), (30) and the above inequality, by induction, we prove the desired statement of Proposition 6.

Proposition 7 *For all $\epsilon > 0$ let*

$$C(n, \epsilon) = \sqrt{\mathbf{E}[(x_n - x'_n)^2; |x_n - c| \leq \epsilon; |x'_n - c| \leq \epsilon]} \quad (33)$$

Suppose that for any $\epsilon > 0$ there exists $N > 0$ such that $C(N, \epsilon) > 0$. Then for any $\epsilon > 0$ there exists $\epsilon_1 > 0$, $M_1 > 0$ and $N_1 > 0$ such that for all $n \geq N_1$

$$C(n, \epsilon_1) > M_1(\lambda - \epsilon)^n \quad (34)$$

Proof: By the averaging property of f and by mean value theorem, for any $\epsilon > 0$ there exists $\epsilon_1 > 0$ such that for all $|x_{n,i} - c| \leq \epsilon_1$ and $|x'_{n,i} - c| \leq \epsilon_1$ where $i = 1, 2, \dots, k$, we have

$$|x_{n+1} - c| \leq \epsilon_1 \text{ and } |x'_{n+1} - c| \leq \epsilon_1$$

and

$$|x_{n+1} - x'_{n+1} - \sum_{i=1}^k \alpha_i (x_{n,i} - x'_{n,i})| \leq \epsilon \sum_{i=1}^k |x_{n,i} - x'_{n,i}|$$

By applying the triangle inequality to the above inequality,

$$\begin{aligned} C(n+1, \epsilon_1) &\geq \sqrt{\mathbf{E}[(x_{n+1} - x'_{n+1})^2; |x_{n,i} - c| \leq \epsilon_1; |x'_{n,i} - c| \leq \epsilon_1, i=1,2,\dots,k]} \\ &\geq \sqrt{\mathbf{E}[(\sum_{i=1}^k \alpha_i (x_{n,i} - x'_{n,i}))^2; |x_{n,i} - c| \leq \epsilon_1, |x'_{n,i} - c| \leq \epsilon_1, i=1,2,\dots,k]} \\ &\quad - \epsilon \sqrt{\mathbf{E}[(\sum_{i=1}^k |x_{n,i} - x'_{n,i}|)^2; |x_{n,i} - c| \leq \epsilon_1, |x'_{n,i} - c| \leq \epsilon_1, i=1,2,\dots,k]} \end{aligned}$$

Hence using the independence and the triangle inequality, we obtain

$$C(n+1, \epsilon_1) \geq \lambda \mathbf{P}[|x_n - c| \leq \epsilon_1]^{k-1} C(n, \epsilon_1) - \epsilon k C(n, \epsilon_1) \quad (35)$$

By weak law of large numbers, there exists $N > 0$ such that for all $n \geq N$,

$$C(n+1, \epsilon_1) \geq (\lambda(1 - \epsilon) - k\epsilon) C(n, \epsilon_1)$$

Since the above is true for any $\epsilon > 0$ (ϵ_1 and N depending on ϵ), Proposition 7 follows immediately by induction.

Proof of Theorem 2: Suppose x_n and f satisfy all assumptions of Theorem 2. We only need to prove that the same sequence x_n satisfies all assumptions of Theorem 1. We only need to check (6). Since f is averaging, $\alpha_i \geq 0$ (by monotonicity) and $\sum_{i=1}^k \alpha_i = 1$. Also $\alpha_i > 0$ at least for two distinct indices (by assumptions of Theorem 2). Hence

$$0 < \lambda = \sqrt{\sum_{i=1}^k \alpha_i^2} < 1$$

For all n , define the essential range of x_n by

$$\mathcal{R}(x_n) = \{u \in [a, b] : \text{for any } \epsilon > 0 \mathbf{P}[|x_n - c| > \epsilon] > 0\}$$

Since $\mathcal{R}(x_0)$ is connected and consists of more than one point, by assumptions of Theorem 2 and by averaging property of f $\mathcal{R}(x_n) = \mathcal{R}(x_0)$ is connected and consists of more than one point. In particular for all n and $\epsilon_1 > 0$ $C(n, \epsilon_1) > 0$. Since $0 < \lambda < 1$, by Proposition 6 and Proposition 7, the condition (6) follows immediately.

Proof of Theorem 3: Since

$$\frac{\hat{x}_n - \mathbf{E}[\hat{x}_n]}{\sqrt{\mathbf{Var}[\hat{x}_n]}} = \frac{x_n - \mathbf{E}[x_n]}{\sqrt{\mathbf{Var}[x_n]}},$$

it is enough to prove the central limit theorem for x_n . Note that f is averaging and since $\mathbf{P}[x_0 > 0] > p_c(\mathbf{G})$, there exists $c > 0$ such that x_n converges to c in probability (see [5]). Moreover f is at least twice continuously differentiable in the neighborhood of $(c, c, \dots, c)$. Since each self-avoiding path connecting two surface sites of $\mathbf{G}$ has length at least 2, there exists $\delta > 0$ such that $\frac{\partial f}{\partial u_i}(c, c, \dots, c) > \delta$ at least two distinct indices i . Hence we only need to

prove (6) to in order apply Theorem 1 for x_n . Let s_t and s_b be the top and bottom sites of $\mathbf{G}$ respectively. For each bond i of $\mathbf{G}$, let $s_1(i)$ and $s_2(i)$ be the two sites of bond i . The variational principle of conductivity says that for any conductance $w_i \geq 0$ of i -th bond of $\mathbf{G}$, we have

$$f(w_1, w_2, \dots, w_k) = C \min_v \sum_{i=1}^k w_i (v(s_1(i)) - v(s_2(i)))^2 \quad (36)$$

where minimum is taken over all real-valued functions v , defined on site set of $\mathbf{G}$, satisfying $v(s_t) = 1$ and $v(s_b) = 0$ and the normalization constant $C > 0$ is chosen to satisfy $f(1, 1, \dots, 1) = 1$ (see [1] for variational principle of conductivity). Let A and B be two bond-disjoint self-avoiding paths connecting two surface sites of $\mathbf{G}$. Let

$$u_i = \begin{cases} 1 & \text{if } i \in A \\ 0 & \text{otherwise} \end{cases}$$

$$u'_i = \begin{cases} 1 & \text{if } i \in B \\ 0 & \text{otherwise} \end{cases}$$

$$\eta_A = f(u_1, u_2, \dots, u_k)$$

$$\eta_B = f(u'_1, u'_2, \dots, u'_k)$$

Then by (36) $\eta_A > 0$ and $\eta_B > 0$. Let $b > a$. For each bond i of $\mathbf{G}$, let

$$w_i = \begin{cases} b & \text{if } i \in A \\ a & \text{otherwise} \end{cases}$$

By applying (36) to w , we get

$$a + (b - a)\eta_A \leq f(w_1, w_2, \dots, w_k) \leq b - (b - a)\eta_B \quad (37)$$

Take any $\epsilon > 0$. Since x_0 is not almost surely a constant, by averaging property of f x_n is not almost surely a constant. Also by weak law of large numbers, there exists N such that

$$\mathbf{P}[|x_N - c| \leq \frac{\epsilon}{2}] > 0$$

For such N , we can choose two distinct real numbers $a_N \in \mathcal{R}(x_N)$ and $b_N \in \mathcal{R}(x_N)$ such that $|a_N - c| < \frac{\epsilon}{2}$. Without loss of generality, let us assume that $a_N < b_N$. For all $n > N$, define a_n and b_n inductively by

$$a_n = f(a_{n-1}, a_{n-1}, \dots, a_{n-1}) = a_{n-1}$$

and

$$b_n = f(w_{n-1,1}, w_{n-1,2}, \dots, w_{n-1,k})$$

where

$$w_{n-1,i} = \begin{cases} b_{n-1} & \text{if } i \in A \\ a_{n-1} & \text{otherwise} \end{cases}$$

Clearly for all $n \geq N$, a_n and b_n are distinct and elements of $\mathcal{R}(x_n)$. Furthermore by (37) $b_n - a_n$ converges to zero when n goes to infinity. Hence for large n , $C(n, \epsilon) > 0$. Since $0 < \lambda < 1$, (6) follows immediately from Proposition 6 and Proposition 7.

References

- [1] Doyle, P. G. and Snell, J. L. (1984) Random walks and electric networks. Math. Assoc. Amer. Washington. DC.
- [2] Durrett, R. (1995) Probability: Theory and Examples. (2nd. ed) Wadsworth
- [3] Griffiths, R. B. and Kaufman, M. (1982) Spin systems on hierarchical lattices. Introduction and thermodynamic limit. Phys. Rev. B. Vol **26** pp. 5022-5032.
- [4] Schlosser, T. and Spohn, H. (1992) Sample-to-Sample Fluctuations in the Conductivity of a Disordered Medium. J. Stat. Phys. **69** pp. 955-967
- [5] Shneiberg, I. (1986) Hierarchical sequences of random variables. Prob. Theor. Appl. **31** pp. 137-141.